

Neoadjuvant - Nasopharyngeal					
PI	CRC	Protocol #/Title	Mechanism	Primary In/Ex Criteria	Status
Dr. Nabar	Natalie Arechiga	NRG-HN001 Randomized Phase II and Phase III Studies of Individualized Treatment for Nasopharyngeal Carcinoma Based on Biomarker Epstein Barr Virus (EBV) Deoxyribonucleic Acid (DNA)	• •	Must have detectable EBV DNA Biopsy proven stage II-IVB nasopharyngeal cancer with no distant metastasis Must not have prior invasive malignancy	Open to accrual
NeoAdjuvant - Squamous Cell Carcinoma of Head and Neck - Oropharyngeal					
PI	CRC	Protocol #/Title	Mechanism	Primary In/Ex Criteria	Status
Dr. Chen	Natalie Arechiga	UCI 22-132: De-escalated Radiation for Human Papillomavirus-Positive Squamous Cell Carcinoma of the Oropharynx	IMRT	Inclusion: Pathologically proven diagnosis of HPV- positive OPSCC. HPV-positivity will be defined as tumors that are p16-positive by immunohistochemistry Clinical stage I, II, or III disease (AJCC Eighth Edition) Exclusion: Patients who have had initial surgical treatment other than the diagnostic biopsy of the primary site or nodal sampling of the neck disease are excluded;	Open to accrual
Adjuvant - Squamous Cell Carcinoma of Head and Neck					
PI	CRC	Protocol #/Title	Mechanism	Primary In/Ex Criteria	Status
Recurrent and/or Metastatic Squamous Cell Carcinoma					
PI	CRC	Protocol #/Title	Mechanism	Primary In/Ex Criteria	Status