

Heme Malignancy Disease-Oriented Team

Clinical Research Treatment Trial Flowchart

Clinical Research Managers:

Stephanie Osorio
Judit Castellanos

Clinical Research Coordinators:

Kelsey McAbee
Harleen Mehrok
Alice Ting
Michael Kunicki
Georgina Alvarez-Diaz

Data Coordinators:

Heather Franson
Neha Ashraf

Newly diagnosed

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

Front Line

ETCTN 10538 (suspended)

Venetoclax+ASTX727 (All oral therapy) for CMML, MDS/MPN with excess blasts

Accrual: 0/5

Coord: Stephanie Osorio
Mechanism: BCL-2 selective inhibitor

UCI 24-121

ASTX030 in Subjects w/Myeloid Neoplasm or in Combo w/Venetoclax in Subjects w/AML or MDS

Accrual: 0/5

Coord: TBD
Mechanism: cytidine deaminase inhibitor

Observational Study

UCI 23-32

Dissecting the mechanism of Interferon Alpha (IFN) response in MPN

Accrual: 13/50

Coord: N/A
Mechanism: observational study

Supportive Care

UCI 20-50

N-Acetylcysteine in MPN to Improve Disease Markers & Symptoms

Accrual 14/27

Coord: Kelsey McAbee
Mechanism: Mucolytic agent (cysteine and GSH precursor)

Newly diagnosed

UCI 24-121

ASTX030 in Subjects
w/Myeloid Neoplasm or in
Combo w/Venetoclax in
Subjects w/AML or MDS

Accrual: 0/5

Coord: TBD
Mechanism: cytidine
deaminase inhibitor

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Low-Risk

Molecularly-Driven

HSCT

High-Risk

Relapsed/Refractory

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

High-Risk

UCI 22-151

LYT-200 in patients w/ R/R AML or high-risk MDS

Accrual: 6/8

Coord: Stephanie Osorio

Mechanism: Galectin-9 monoclonal antibody

UCI 23-113

Oral GLB-001 in patients w/ R/R AML or high-risk MDS

Accrual: 2/7

Coord: Stephanie Osorio

Mechanism: Selective molecular glue degrader

Low-Risk

UCI 21-239

IRAK 1/4 inhibitor, R289, in patients w/ refractory or resistant lower-risk MDS

Accrual: 1/5

Coord: Stephanie Osorio

Mechanism: IRAK1/4 inhibitor

Molecularly-Driven

Newly diagnosed

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

Intensive

ETCTN-10596 (SUSPENDED)

SNDX-5613 + Daunorubicin
 and Cytarabine in Newly
 Diagnosed Acute Myeloid
 Leukemia (NPM1
 Mutated/FLT3 Wildtype with
 Higher-Risk Features or
 MLL/KMT2A Rearranged)

Accrual: 0/5

Coord: Stephanie Osorio
 Mechanism: menin inhibitor

Non-Intensive

ETCTN-10630

Ladademstat in Combination with
 Venetoclax and Azacitidine in
 Patients with Post MDS
 Transformation to AML

Accrual: 2/7

Coord: Stephanie Osorio
 Mechanism: LSD1 inhibitor

UCI 24-121

ASTX030 in Subjects w/Myeloid
 Neoplasm or in Combo
 w/Venetoclax in Subjects
 w/AML or MDS

Accrual: 0/5

Coord: TBD
 Mechanism: cytidine
 deaminase inhibitor

KMT2A-r/NPM1-m**UCI 23-44**

Venetoclax/Azacitidine v.s
 Venetoclax+ KO-530 v.s
 cytarabine/daunorubicin
 (7+3)+ KO-539 in AML

Accrual: 8/10

Coord: Stephanie Osorio
 Mechanism: menin
 inhibitor

FLT3 mutation**UCI 21-216**

Gilteritinib+Venetoclax+Azacitidine in patients w/ FLT3 mutant AML not eligible for intensive induction chemotherapy

Accrual: 2/5

Coord: Stephanie Osorio
 Mechanism: FLT3 inhibitor

Newly diagnosed

KMT2A-r/NPM1-m

UCI 25-20

Assessing Ziftomenib in
Combination with Either
Standard of Care Nonintensive
(Venetoclax+Azacitidine) or
Intensive (7+3) Therapy in
Patients with Untreated NPM1
Mutated or KMT2A
Rearranged AML

Accrual: 0/5

Coord: TBD

Mechanism: menin inhibitor

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Relapsed/Refractory

2nd Line+

UCI 23-113

Oral GLB-001 in patients w/
R/R AML or high-risk MDS

Accrual: 2/7

Coord: Stephanie Osorio
Mechanism: Selective
molecular glue degrader

UCI 22-151

LYT-200 in patients w/ R/R
AML or high-risk MDS

Accrual: 6/8

Coord: Stephanie Osorio
Mechanism: Galectin-9
monoclonal antibody

UCI 22-81

HM43239 in patients w/ R/R
AML

Accrual: 1/6

Coord: Stephanie Osorio
Mechanism: FLT3 inhibitor

UCI 24-48

DFP-10917+Venetoclax in
R/R AML

Accrual: 4/5

Coord: Stephanie Osorio
Mechanism: Deoxycytidine
nucleoside analogue (DNA
synthesis inhibitor)

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Molecularly-Driven

≤1 Line Only

UCI 25-105

LYT-200 in patients w/ R/R
AML or high-risk MDS

Accrual: 6/8

Coord: Stephanie Osorio
Mechanism: Galectin-9
monoclonal antibody

Relapsed/Refractory

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

Molecularly-Driven

KMT2A-r/NPM1-m

UCI 23-44

Venetoclax/Azacitidine v.s
Venetoclax+ KO-530 v.s
cytarabine/daunorubicin (7+3)+
KO-539 in AML

Accrual: 8/10

Coord: Stephanie Osorio
Mechanism: menin inhibitor

UCI 23-154

Ziftomenib combinations for
the KMT2A-rearranged/NPM1
mutant R/R AML

Accrual: 3/5

Coord: Stephanie Osorio
Mechanism: menin inhibitor

UCI 25-24

8-Chloro-Adenosine in
Combination with Decitabine and
Venetoclax in Patients with R/R
AML

Accrual: 0/5

Coord: TBD
Mechanism: RNA directed
nucleoside

Maintenance

High-Risk, HSCT

Salvage Therapy

Newly diagnosed

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

Ph+ only

EA9181

Steroids +TKI w/
chemotherapy or
Blinatumomab for BCR-ABL
positive adult patients

Accrual 14/35

Coord: Judit Castellanos
Mechanism: BiTE binding to
CD19 (on B-cell) and CD3 (on
T-cells) and PD-1 inhibitor

SWOG-21CTP-LEUK01

Phase II Trial of Asciminib,
Dasatinib, Prednisone, and
Blinatumomab for Participants
with Newly Diagnosed
Philadelphia Chromosome
Positive (PH+) Acute
Lymphoblastic Leukemia
Accrual: 0/5

Coord: Stephanie Osorio/Judit
Castellanos
Mechanism:

Ph- only

Age 22-55 years & BMI <35kg/m2

UCI 22-125 (closed to accrual)

Calaspargase pegol for tx of
adults 22-55y/o w/ newly
diagnosed Ph- ALL

Accrual: 0/5

Coord: Judit Castellanos
Mechanism: PEGylated
conjugate L-asparaginase

Age ≥ 18 years & < 40 years,
CD22+ (≥ 20%)

A041501 (Suspended)

Addition of Inotuzumab
Ozogamicin to frontline
therapy in young adults (18-
39y/o)

Accrual: 10/15

Coord: Judit Castellanos
Mechanism: conjugated anti-
CD22 monoclonal antibody

Age 5 to <30 years & High Risk ALL

UCI 21-14

Levocarnitine for Asparaginase
hepatotoxicity in ALL patients

Accrual: 0/5 (opened 11/3/23)

Coord: Judit Castellanos
Mechanism: Oxidative stress
reducer & inflammatory
modulator

UCI 25-104

subcutaneous blinatumomab
vs. intravenous blinatumomab
in newly diagnosed adults with
(Ph)- B cell precursor ALL

Accrual: 0/5

Coord: TBD
Mechanism: Bispecific T-cell
engager (BiTE)

Observational

UCI 21-236

Addressing the Hispanic
Cancer Disparity in B Cell
Acute Lymphoblastic
Leukemia
Accrual: NA

Coord: NA
Mechanism: Observational

Relapsed/Refractory

■ Open to Accrual

■ Low Accruing

■ Pending Activation/Suspended

Molecularly-Driven

CR w/ MRD+

CD22+ ($\geq 20\%$)**A041703 (SUSPENDED)**

Inotuzumab Ozogamicin followed by
Blinatumomab for ph- CD22-positive
newly diagnosed or R/R ALL patients

Accrual: 2/5 (only open for R/R)

Coord: Judit Castellanos
Mechanism: antibody-drug
conjugate combining a monoclonal
antibody targeting CD22 on B-
lymphoblast with the cytotoxic agents

Newly diagnosed



Open to Accrual



Low Accruing



Pending Activation/Suspended

High-Risk

S1925

Venetoclax+Obnutumab early intervention vs. delayed therapy in asymptomatic high-risk CLL/SLL

Accrual: 4/10

Coord: Kelsey McAbee
Mechanism: BCL2 inhibitor +anti-CD20 monoclonal antibody

Front Line

UCI 23-156

Sonrotoclax (BGB-11417) + Zanubrutinib (BGB-3111) v.s. Venetoclax +Obinutuzumab

Accrual: 3/7

Coord: Kelsey McAbee
Mechanism: BTK + BCL2 inhibition

Relapsed/Refractory

■ Open to Accrual

■ Low Accruing

■ Pending Activation/Suspended

3rd Line+**UCI 22-134**

Oral AS-1763 in patients w/
previously treated CLL/SLL or NHL

Accrual: 4/9

Coord: Kelsey McAbee
Mechanism: BTK inhibitor for both
wild-typ and C481S-mutant type

UCI 24-12

Study to Evaluate the BTK Degradar,
ABBV-101, in Participants With B-cell
Malignancies

Accrual: 1/5

Coord: Kelsey McAbee
Mechanism: BTK inhibitor/f ABBV-
101 monotherapy

2nd Line+**UCI 24-190**

A Phase III Study of Sonrotoclax Plus
Anti-CD20 Antibody Therapies Versus
Venetoclax + Rituximab in Patients With
R/R CLL/SLL

Accrual: 0/7

Coord: Stephanie Osorio/Judit
Castellanos
Mechanism: Anti-CD20 Antibody

UCI 24-32

Epcoritamab as Consolidation Therapy
for High-risk Patients with Chronic
Lymphocytic Leukemia on Bruton
Tyrosine Kinase Inhibitor

Accrual: 0/24

Coord: Stephanie Osorio/Judit
Castellanos
Mechanism: BTKi/Epcoritamab

UCI 25-04

A Phase I/II, multicenter study
evaluating point-of-care manufactured
GLPG5101 (19CP02) in subjects with
R/R B-cell non-Hodgkin lymphoma
(CP0201NHL)

Accrual: 0/5

Coord: TBD
Mechanism: FMC63 anti-CD19 CAR T

UCI 25-148

IgPro20 in Subjects with Chronic
Lymphocytic Leukemia / Small
Lymphocytic Lymphoma and
Secondary Hypogammaglobulinemia
Experiencing Severe or Recurrent
Infections

Accrual: 0/5

Coord: TBD
Mechanism: protein liquid preparation
of polyvalent human IgG

Relapsed/Refractory

2nd Line+

UCI 23-167

Phase I- TERN-701 in patients
w/CML

Accrual: 2/5

Coord: Kelsey McAbee
Mechanism: STAMP inhibitor

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Newly Diagnosed

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Post ASCT

Front Line

Bispecific

UCI 23-158 (SUSPENDED)

Phase I/II Study of Linvoseltamab (Anti-BCMA X Anti-CD3 Bispecific Antibody) in Previously Untreated Patients with Symptomatic Multiple Myeloma

Accrual: 1/6 (opened 3/29/24)

Coord: Alice Ting
Mechanism: Bispecific antibody (BCMA x CD3)

High-Risk

ETCTN 10612

A Randomized Phase 2 Study of Daratumumab-Selinexor-Velcade-Dexamethasone (Dara-SVD) for High-Risk Newly Diagnosed Multiple Myeloma

Accrual: 7/8 (opened 4/25/24)

Coord: Alice Ting
Mechanism: selective inhibitor of nuclear export

UCI 23-49

ELRANATAMAB (PF-06863135) + DARATUMUMAB + LENALIDOMIDE VERSUS DARATUMUMAB + LENALIDOMIDE + DEXAMETHASONE IN TRANSPLANT-INELIGIBLE PARTICIPANTS WITH NEWLY-DIAGNOSED MULTIPLE MYELOMA
Accrual: 0/5

Coord: TBD
Mechanism: IgG1 kappa (IgG1κ) MAB

Relapsed/Refractory

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Molecularly-Driven

2nd Line+UCI 22-190

Teclistamab monotherapy vs.
PVD or KD in patients received
1-3 prior lines of therapy

Accrual: 3/6

Coord: Alice Ting
Mechanism: CD3 x BCMA BiTE

3rd Line+

CAR-T

UCI 24-02 (SUSPENDED)

Descartes-15 in R/R MM
Accrual: 3/5

Coord: Mike Kunicki
Mechanism: CAR-T, BCMA

ALLIANCE-A062102

Iberdomide Maintenance
Therapy Following Idecabtagene
Vicleucel CAR-T in R/R MM
Accrual: 0/5

Coord: Judit Castellanos
Mechanism: cereblon (CRBN)
modulating agent

Maintenance

Relapsed/Refractory**UCI 25-62**

EXPANDED ACCESS PROTOCOL
(EAP) FOR SUBJECTS
RECEIVING IDECABTAGENE
VICLEUCEL THAT IS
NONCONFORMING FOR
COMMERCIAL RELEASE

Accrual: 0/5

Coord: TBD

Mechanism: CAR-T

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Molecularly-Driven

2nd Line+

Maintenance

3rd Line+

Newly diagnosed

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

Front Line

UCI 23-17

Odronextamab (REGN1979) vs.
investigator's choice in patient w/ FL

Accrual: 0/5 (3/20/24)

Coord: Harleen Mehrok

Mechanism: Anti-CD20 x Anti-CD3
bispecific antibody

SWOG 2308

MOSUNETUZUMAB VS. RITUXIMAB
FOR LOW TUMOR BURDEN
FOLLICULAR LYMPHOMA

Accrual: 0/5

Coord: Stephanie Osorio/Judit
Castellanos

Mechanism: Anti-CD20 IgG1 kappa
antibody

Relapsed/Refractory

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

Cell Therapy

UCI 25-04

A Phase I/II, multicenter study evaluating point-of-care manufactured GLPG5101 (19CP02) in subjects with R/R B-cell non-Hodgkin lymphoma (CP0201NHL)

Accrual: 0/5

Coord: TBD

Mechanism: FMC63 anti-CD19 CAR T

UCI 23-114

Safety and Efficacy of IMPT-314, a CD19/20 Bispecific Chimeric Antigen Receptor (CAR) T Cell Therapy in B-cell NHL

Accrual: 4/7

Coord: Judit Castellanos

Mechanism: CD19/20 bispecific CAR

3rd Line+

UCI 22-134

Oral AS-1763 in patients w/ previously treated CLL/SLL or NHL

Accrual: 4/9

Coord: Kelsey McAbee

Mechanism: BTK inhibitor for both wild-typ and C481S-mutant type

Consolidation

S2114

Consolidation therapy following CD19 CAR T-cell tx

Accrual: 0/6

Coord: Michael Kunicki

Mechanism: bite/mab

Molecularly-Driven

Outpatient

Relapsed/Refractory

2+ Lines

UCI 24-12

Study to Evaluate the BTK Degradar, ABBV-101, in Participants With B-cell Malignancies

Accrual: 1/5

Coord: Kelsey McAbee

Mechanism: BTK inhibitor/f ABBV-101 monotherapy

UCI 25-53

JNJ-90014496, an Autologous CD19/CD20 Bi-specific CAR-T Cell Therapy in Adult Participants with B-cell Non-Hodgkin Lymphoma

Accrual: 0/5

Coord: TBD

Mechanism: Bi-specific CAR T

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Cell Therapy

Relapsed/Refractory

■ Open to Accrual

■ Low Accruing

■ Pending Activation/Suspended

EBV+

Molecularly-Driven

2 + Lines

UCI 24-12

Study to Evaluate the BTK Degradar, ABBV-101, in Participants With B-cell Malignancies

Accrual: 1/5

Coord: Kelsey McAbee
Mechanism: BTK inhibitor/f ABBV-101 monotherapy

UCI 25-04

A Phase I/II, multicenter study evaluating point-of-care manufactured GLPG5101 (19CP02) in subjects with R/R B-cell non-Hodgkin lymphoma (CP0201-NHL)

Accrual: 0/5

Coord: TBD
Mechanism: FMC63 anti-CD19 CAR T

3rd Line+UCI 22-134

Oral AS-1763 in patients w/ previously treated CLL/SLL or NHL

Accrual: 4/9

Coord: Kelsey McAbee
Mechanism: BTK inhibitor for both wild-typ and C481S-mutant type

Relapsed/Refractory

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

Cell Therapy

Molecularly-Driven

3rd Line+

UCI 22-134

Oral AS-1763 in patients w/
previously treated CLL/SLL or NHL

Accrual: 4/9

Coord: Kelsey McAbee
Mechanism: BTK inhibitor for both
wild-typ and C481S-mutant type

2+ Lines

UCI 24-12

Study to Evaluate the BTK
Degradar, ABBV-101, in
Participants With B-cell
Malignancies

Accrual: 1/5

Coord: Kelsey McAbee
Mechanism: BTK inhibitor/f ABBV-
101 monotherapy

UCI 25-04

A Phase I/II, multicenter study
evaluating point-of-care
manufactured GLPG5101 (19CP02)
in subjects with R/R B-cell non-
Hodgkin lymphoma (CP0201NHL)

Accrual: 0/5

Coord: TBD
Mechanism: FMC63 anti-CD19 CAR
T

Newly diagnosed

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

75 y/o Older

S1918 (SUSPENDED)

R-miniCHOP w/ or w/o oral
Azacitidine in patients 75 y/o or
older

Accrual: 5/10

Coord: Regan Dagenhart
Mechanism: Oral
hypomethylating agent

Relapsed/Refractory

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Primary Relapsed/Refractory

Cell Therapy- CRS mgmt

UCI 23-193 (IRB initial approval)
CTO1681 for the Prevention and
Treatment of CRS in Patients with
DLBCL receiving Chimeric Antigen
Receptor T-Cell Therapy
Accrual: 1/5

Coord: Judit Castellanos
Mechanism: PGE2 & PGI2 agonist

Secondary Relapsed/Refractory

UCI 20-126

CB-010, CRISPR-edited
allogeneic anti-CD19 CAR-T
cell therapy

Accrual: 7/10

Coord: Michael K.
Mechanism: anti-CD19

**CHIMERIC ANTIGEN
RECEPTOR**

Outpatient

Relapsed/Refractory

■ Open to Accrual

■ Low Accruing

■ Pending Activation/Suspended

Molecularly-Driven

Tertiary Relapsed/Refractory

S2114Consolidation therapy
following CD19 CAR T-cell tx

Accrual: 0/6

Coord: Michael Kunicki
Mechanism: bite/mab

2+ Lines

UCI 24-12Study to Evaluate the BTK Degradar,
ABBV-101, in Participants With B-cell
Malignancies

Accrual: 1/5

Coord: Kelsey McAbee
Mechanism: BTK inhibitor/f ABBV-101
monotherapy**UCI 25-53**JNJ-90014496, an Autologous
CD19/CD20 Bi-specific CAR-T Cell
Therapy in Adult Participants with B-cell
Non-Hodgkin Lymphoma

Accrual: 0/5

Coord: TBD
Mechanism: Bi-specific CAR T**UCI 25-04**A Phase I/II, multicenter study
evaluating point-of-care manufactured
GLPG5101 (19CP02) in subjects with
R/R B-cell non-Hodgkin lymphoma
(CP0201NHL)

Accrual: 0/5

Coord: TBD
Mechanism: FMC63 anti-CD19 CAR T**UCI 23-114**Safety and Efficacy of IMPT-314, a
CD19/20 Bispecific Chimeric Antigen
Receptor (CAR) T Cell Therapy in B-
cell NHL

Accrual: 4/7

Coord: Judit Castellanos
Mechanism: CD19/20 bispecific CAR

Newly Diagnosed

COG-AHOD2131

Standard Therapy with
Immuno-oncology Therapy
for Newly Diagnosed Stage I
and II Classical Hodgkin
Lymphoma

Accrual: 1/5

Coord: Judit Castellanos/
Stephanie Osorio
Mechanism:

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Molecularly-Driven

Basket study

Relapsed/Refractory

 ≥ 1 Line

UCI 25-21

Pembrolizumab and GVD with
ctDNA-guided Consolidation
in Patients with Relapsed or
Refractory Classic Hodgkin
Lymphoma

Accrual: 1/5

Coord: TBD

Mechanism: PD-1 inhibitor

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Molecularly-Driven

Basket study

Newly diagnosed

COG ANHL1931

Nivolumab + chemo-
immunotherapy

Accrual: 2/5

Coord: Regan Dagenhart

Mechanism: PD1 inhibitor

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Relapsed/Refractory

■ Open to Accrual

■ Low Accruing

■ Pending Activation/Suspended

Molecularly-Driven

Consolidation

S2114

Consolidation therapy following
CD19 CAR T-cell tx

Accrual: 0/6

Coord: Michael Kunicki
Mechanism: bite/mab

2+ Lines

UCI 24-12

Study to Evaluate the BTK
Degradar, ABBV-101, in
Participants With B-cell
Malignancies

Accrual: 1/5

Coord: Kelsey McAbee
Mechanism: BTK inhibitor/f ABBV-
101 monotherapy

Relapsed/Refractory

■ Open to Accrual ■ Low Accruing ■ Pending Activation/Suspended

Molecularly-Driven

Cell Therapy

UCI 23-114

Safety and Efficacy of IMPT-314, a CD19/20 Bispecific Chimeric Antigen Receptor (CAR) T Cell Therapy in B-cell NHL
Accrual: 4/7

Coord: Judit Castellanos
Mechanism: CD19/20 bispecific CAR

UCI 25-04

A Phase I/II, multicenter study evaluating point-of-care manufactured GLPG5101 (19CP02) in subjects with R/R B-cell non-Hodgkin lymphoma (CP0201NHL)

Accrual: 0/5

Coord: TBD
Mechanism: FMC63 anti-CD19 CAR T

2+ Lines

UCI 24-12

Study to Evaluate the BTK Degradar, ABBV-101, in Participants With B-cell Malignancies

Accrual: 1/5

Coord: Kelsey McAbee
Mechanism: BTK inhibitor/f ABBV-101 monotherapy

3rd line+

UCI 22-134

Oral AS-1763 in patients w/ previously treated CLL/SLL or NHL

Accrual: 4/9

Coord: Kelsey McAbee
Mechanism: BTK inhibitor for both wild-typ and C481S-mutant type

Relapsed/Refractory

■ Open to Accrual
 ■ Low Accruing
 ■ Pending Activation/Suspended

2nd Line+

UCI 21-99

ONO-4685 given as monotherapy

Accrual: 4/10

Coord: TBD

Mechanism: CD3-bispecific antibody targeting PD-1

UCI 25-74

A Multicenter, Open-Label, First-In-Human, Multiple Expansion Cohort, Phase I/II Study to Evaluate the Safety and Efficacy of DR-01 in Subjects with LGLL or Cytotoxic Lymphomas

Accrual: 0/5

Coord: TBD

Mechanism: non fucosylated human IgG1 anti CD94 antibody

UCI 24-188

PHASE 2 STUDY OF PTX-100 MONOTHERAPY IN PATIENTS WITH R/R CUTANEOUS T-CELL LYMPHOMA

Accrual: 0/5

Coord: TBD

Mechanism: GGTase 1 inhibitor

UCI 25-142

Study of KK2223 in Participants with Relapsed or Refractory T-cell Non-Hodgkin Lymphoma

Accrual: 0/5

Coord: TBD

Mechanism: FMC63 anti-CD19 CAR T

Relapsed/Refractory

ECOG EA4232

Phase III Study to Evaluate Autologous Stem Cell Transplant in Patients with Peripheral T Cell Lymphoma that Achieved a First Complete Remission (CR1) Following Induction Therapy (PTCL-STAT)

Accrual: 0/5

Coord: TBD

Mechanism: ASCT

3rd Line+

UCI 21-99

ONO-4685 given as monotherapy

Accrual: 4/10

Coord: TBD

Mechanism: CD3-bispecific antibody targeting PD-1

1st Line+

UCI 25-142

Study of KK2223 in Participants with Relapsed or Refractory T-cell Non-Hodgkin Lymphoma

Accrual: 0/5

Coord: TBD

Mechanism: naked CD4 antibody

Newly Diagnosed

UCI 22-18

Investigation of MRD in patients with treatment-naïve systemic T cell lymphoma treated with a brentuximab-containing regimen

Accrual: 0/5

Coord: TBD

Mechanism:

Allo-SCT Conditioning

UCI 21-90

Risk-ADAPTEd conditioning regimen
for AHSCT

Accrual: 23/48

Coord: Heme CRCs

Allo-SCT Supportive Care

UCI 22-188

Prospective evaluation of CMV-TCIP
directed Letemovir ppx after AHCT

Accrual: 11/50

Coord: Heme CRCs

Auto-SCT Maintenance

CAR-T**UCI 20-126**

CB-010, CRISPR-edited allogeneic anti-CD19 CAR-T cell therapy

Accrual: 7/10

Coord: Michael K.

Mechanism: anti-CD19 CAR-T

UCI 23-114

Safety & Efficacy of IMPT-314, a CD19/20 Bispecific CAR-T in Participants with R/R B-Cell NHL

Accrual: 4/7

Coord: Judit Castellanos

Mechanism: CD19/20 bispecific CAR

UCI 24-02

Descartes-15 in R/R MM

Accrual: 3/5

Coord: Mike K.

Mechanism: CAR-T, BCMA

Supportive Care**UCI 23-193**

CTO1681 for the Prevention and Treatment of CRS in Patients with DLBCL receiving CAR-T Therapy

Accrual: 1/5

Coord: Alice Ting

Mechanism: PGE2 & PGI2 agonist

Post CAR-T**S2114**

Consolidation Therapy Following CD19 CAR-T for R/R Large B-cell Lymphoma or Grade IIIB Follicular Lymphoma

Accrual: 0/6

Coord: Michael Kunicki

Mechanism: BiTE/mAb

Alliance-A062102

Iberdomide Maintenance Therapy Following Ide-Cel CAR-T in R/R Multiple Myeloma

Accrual: 0/5

Coord: TBD

Mechanism: Cereblon (CRBN) modulating agent



Supportive Care

Long-Term FU

UCI 14-03
Role of Inflammation in the
Pathogenesis of
Myeloproliferative Neoplasm

UCI 21-184
Long-term safety of CAR-T
inpatient w/ heme malignancies

Accrual: 4/5
Coord: Miranda Duron

UCI 15-65

Effect of candidate blood
cancer therapies on normal
human lymphocytes

UCI 24-31
Long-Term Follow-up Protocol for
Subjects Treated With Gene-
Modified T Cells

Accrual: 0/5
Coord: TBD

UCI 25-126
A long-term follow-up study for
patients treated with Galapagos
CAR T-cell therapies

Accrual: 0/5
Coord: TBD
Mechanism: N/A



HLH

UCI 23-189

Frontline Ruxolitinib with De-Intensified HLH-94 for Adults with Hemophagocytic Lymphohistiocytosis (HLH)

Accrual: 0/5

Coord: Stephanie Osorio

UCI 25-74

A Multicenter, Open-Label, First-In-Human, Multiple Expansion Cohort, Phase I/II Study to Evaluate the Safety and Efficacy of DR-01 in Subjects with LGLL or Cytotoxic Lymphomas

Accrual: 0/5

Coord: TBD

Mechanism: non fucosylated human IgG1 anti CD94 antibody

UCI 25-04

A Phase I/II, multicenter study evaluating point-of-care manufactured GLPG5101 (19CP02) in subjects with R/R B-cell non-Hodgkin lymphoma (CP0201NHL)

Accrual: 0/5

Coord: TBD

Mechanism: FMC63 anti-CD19 CAR T

UCI 25-63

EXPANDED ACCESS PROTOCOL (EAP) FOR SUBJECTS RECEIVING LISOCABTAGENE MARALEUCEL THAT IS NONCONFORMING FOR COMMERCIAL RELEASE

Accrual: 0/5

Coord: TBD

Mechanism: anti-CD19 CAR T

CAR-T

UCI 24-152

NXC-201 for the Treatment of Patients with Relapsed or Refractory AL Amyloidosis

Accrual: 0/10

Coord: TBD

Mechanism: anti-CD3 CAR T